

Cotoneaster 'Emerald Beauty'

Key Features

- ✓ Improved fire blight resistance.
- ✓ Improved branching.
- ✓ Fuller plant with fewer touches.



Oregon State University
College of Agricultural
Sciences

Characteristics

Genus & Species:	<i>Cotoneaster x suecicus</i> (Code: OSUCOT2)
Protection Status:	PPAF
USDA Zone:	USDA Zone 5 (-20°F or -28.9°C). Currently Testing
Bloom:	White blooms in spring followed by colorful orange fall berries.
Bloom Time:	Spring-summer.
Foliage:	Shiny small dark green foliage. Flush purple with cool fall temperatures.
Habit:	Well-branched, compact spreading groundcover. 2' x 5'.
Propagation:	Softwood cuttings. <i>License required for propagation.</i>
Culture:	Happy in sun to part shade in average soils.
Additional Comments:	Exhibiting improved fire blight resistance.
Uses:	Container specimens, garden specimen, mass landscape plantings.

Approximate Finishing Time

1 Gallon:	10-12 months
2 Gallon:	16-18 months

Comparison: C. 'Coral Beauty'	Key Differences:	<ul style="list-style-type: none"> ✓ 'Emerald Beauty' improved fire blight resistance vs. less resistant. ✓ 'Emerald Beauty' improved branching vs. less branching. ✓ 'Emerald Beauty' compact denser habit, 2'x 5' vs. more open habit 3' x 6'.
--------------------------------------	-------------------------	--

PlantHaven Disclaimer: These Grower Notes are based on our experience with this crop. Growth will vary depending on location, climate, growing medium and cultural practices. Where fertilizer & chemical use is involved, read the labels, follow the manufacturer's instructions & practice careful trialing methods.

Note: All PlantHaven varieties are subject to patent protection: unlicensed propagation prohibited.

